

ORIGINAL ARTICLE

Factors Associated with Dog Rabies Vaccination in Bohol, Philippines: Results of a Cross-Sectional Cluster Survey Conducted Following the Island-Wide Rabies Elimination Campaign

S. Davlin^{1,2}, S. M. Lapiz³, M. E. Miranda⁴ and K. Murray⁵

¹ School of Public Health, The University of Texas Health Science Center at Houston, Houston, TX, USA

² Department of Obstetrics and Gynecology, Center for Interdisciplinary Research in Women's Health, The University of Texas Medical Branch at Galveston, Galveston, TX, USA

³ Office of the Provincial Veterinarian, Tagbilaran City, Province of Bohol, Philippines

⁴ Global Alliance for Rabies Control, Asia Office, Laguna, Philippines

⁵ Department of Pediatrics, National School of Tropical Medicine, Baylor College of Medicine, Houston, TX, USA

Impacts

- The Visayas region of the Philippines has introduced new rabies control programmes in recent years; therefore, studies on dog populations and factors associated with rabies vaccination are vital.
- Dog ownership is common in Bohol with a human-to-dog ratio of 4 : 1, yielding an estimated owned-dog population of almost 300 000.
- Overall, 71% of dogs were reported vaccinated at some point in their lives. Age was significantly associated with vaccination. Dogs aged 12–23 months had 4.6 times greater odds of vaccination than those aged 3–11 months ($P = 0.002$).

Keywords:

Rabies; dog; cross-sectional; Philippines; cluster survey

Correspondence:

S. Davlin. Department of Obstetrics and Gynecology, Center for Interdisciplinary Research in Women's Health, The University of Texas Medical Branch at Galveston, 301 University Boulevard, Galveston, TX 77555-0587, USA. Tel.: 409 772 3986; Fax: 409 747 5129; E-mail: sldavlin@utmb.edu

Received for publication May 14, 2012

doi: 10.1111/zph.12026

Summary

The Philippines has a long history of rabies control efforts in their dog populations; however, long-term success of such programmes and the goal of rabies elimination have not yet been realized. The Bohol Rabies Prevention and Elimination Program was developed as an innovative approach to canine rabies control in 2007. The objective of this study was to assess canine rabies vaccination coverage in the owned-dog population in Bohol and to describe factors associated with rabies vaccination 2 years after implementation of the programme. We utilized a cross-sectional cluster survey based on the World Health Organization's Expanded Programme on Immunization coverage survey technique. We sampled 460 households and collected data on 539 dogs residing within these households. Seventy-seven per cent of surveyed households reported owning at least one dog. The human-to-dog ratio was approximately 4 : 1, and the mean number of dogs owned per household was 1.6. Based on this ratio, we calculated an owned-dog population of almost 300 000. Overall, 71% of dogs were reported as having been vaccinated for rabies at some time in their lives; however, only 64% of dogs were reported as having been recently vaccinated. Dogs in our study were young (median age = 24 months). The odds of vaccination increased with increasing age. Dogs aged 12–23 months had 4.6 times the odds of vaccination compared to dogs aged 3–11 months (95% CI 1.8–12.0; $P = 0.002$). Confinement of the dog both day and night was also associated with increased odds of vaccination (OR = 2.1; 95% CI 0.9–4.9; $P = 0.07$), and this result approached statistical significance. While the programme is on track to meet its goal of 80% vaccination coverage, educational efforts should focus on the need to confine dogs and vaccinate young dogs.

Introduction

Rabies is an ancient disease. Laws governing rabid dogs have been found as far back as 2300 BC (Baer, 2007). It is plausible that rabies has been a problem in humans since the dog was first domesticated around 14 000 years ago (Rupprecht et al., 2008). Exposure to rabid dogs accounts for 99% or more of all human cases (WHO, 1992), and 99% of those human cases arise in the 80+ countries where canine rabies is still endemic (WHO, 2005a). The present canine rabies problem is unique to the developing world, and it is estimated that 55 000 people die from rabies in Africa and Asia every year (Knobel et al., 2005). Most developed countries have either eliminated rabies completely or reduced it to wildlife reservoirs (WHO, 2005a). With an estimated 500 million dogs worldwide (Macpherson et al., 2000), and half the world's human population living in areas where canine rabies is a real threat (WHO, 2005a), effective canine rabies control programmes are essential to controlling the disease in humans. Both empirical and statistical evidence suggests that maintaining rabies vaccination coverage of 70% in the dog population is effective in controlling the disease in dogs (Coleman and Dye, 1996).

Rabies affects Asia disproportionately. At least 31 000 human deaths due to rabies occur in Asia every year (Knobel et al., 2005). Yet, successful models for rabies elimination exist in Asia. Japan, Taiwan and Peninsular Malaysia have all eliminated endemic rabies in dogs (WHO, 2002). In the Philippines, epidemiological studies on rabies in both humans and animals began as early as the 1960s, and compulsory vaccination of dogs was instituted in Manila during the 1950s (Fishbein et al., 1991). In 2007, the Republic of the Philippines enacted RA 9482, better known as, 'The Rabies Act of 2007', with the goal of national elimination of human rabies by 2020 (Republic of the Philippines, 2011).

The Visayas region of the Philippines has been the target of rabies control programmes in recent years due to the high incidence of the disease in this area. Specifically, the Province of Bohol introduced the Rabies Prevention and Eradication Program beginning in 2007 (Dodet, 2010). Major components of this programme include vaccination of at least 80% of the dog population and to eliminate stray dogs completely. The first island-wide vaccination campaign was conducted in 2007 using *Rabisin*[®] (Merial Animal Health, Lyon, France), a vaccine offering 2-year protection. The campaign has continued annually thereafter. As a result of the programme, only one human rabies case has been reported in Bohol since 2008 (Global Alliance for Rabies Control, 2011).

The aims of this study were to describe the dog population in Bohol and to examine what factors are associ-

ated with rabies vaccination. As Bohol is the subject of an ongoing rabies control programme, it is our hope that both programme evaluation and dog population studies will be conducted on an annual basis and that this baseline data will help determine the direction of future studies.

Methods

Study population

This study was conducted in Bohol, Philippines. The Province of Bohol consists of 48 municipalities. A barangay is the smallest governmental unit in the Philippines (Green et al., 2002). Within each municipality, there are between 8 (Corella) and 67 (Loon) barangays for a total of 1109 barangays on the island. The smallest barangay is Tiwi, in the municipality of Loon, with a population of 84. The largest barangay is Cogon, in the municipality of Tagbilaran City, with a population of 17 266. The total population of Bohol is 1 230 110 (2007 census data).

Data collection

The data for this study came from a cross-sectional survey administered by the Office of the Provincial Veterinarian (OPV) on the island of Bohol during June and July of 2009. We utilized a modified version of multistage cluster sampling technique with population proportional to size (PPS) sampling with replacement that was based on the World Health Organization's Expanded Programme on Immunization (EPI) coverage survey technique (Henderson and Sundaresan, 1982; Lemeshow and Robinson, 1985; Bennett et al., 1991; WHO, 2005b). Based on *a priori* evidence, we assumed a design effect of 1.18. We determined a sample size of 460 households, consisting of 46 clusters (barangays) and 10 households per cluster, was both adequate and feasible to estimate rabies vaccination coverage in the dog population.

In the first stage of data collection, we enumerated the population size of each barangay based on census data and arranged them in alphabetical order. The sampling interval was determined by dividing the total population of Bohol (1 230 110) by the number of clusters we wished to collect (46). The first cluster (barangay) was selected using a randomly generated five-digit number and matching it to the first barangay in our list with a cumulative population greater than or equal to the random number. Cluster 2 was identified by adding the sampling interval to the random number. Subsequent clusters were selected by adding the sampling interval to the previously generated number until a total of 46 clusters were identified. Clusters that were located within the same municipality were retained, and when the end of the list

was reached before all 46 clusters were selected, then selection continued at the top of the list.

Once all 46 clusters were identified, selection of the household was carried out using simple random sampling in the field. The basic sampling unit was the individual household, and a household was defined as those individuals who shared a kitchen. Households were typically selected using a list obtained in the barangay of all the households. If this was not available, we obtained a list of all the streets. Numbers were assigned to each household or street, and corresponding numbers were placed in a hat. One member of the field team drew a number from the hat. The correspondingly numbered household or street was the starting point of the survey. Only the street and/or the first household was randomly selected. Each subsequent household was chosen by going to the next closest front door of the previous household. If no one was at home in the selected household, then the next closest front door to that house was selected. Households were selected in this manner until a total of 10 household questionnaires were obtained in each cluster.

The survey consisted of two distinct questionnaires, and it was administered to any household member age 15 or older who agreed to participate. Questionnaires were administered during the daytime, and households were not revisited. Interviews were conducted either in English or in the local dialect. The first questionnaire was designed to collect household demographics such as interviewee age, sex, relationship to head of household, total number of household members, number of children <15 years in the household and whether the head of household was employed. We also assessed knowledge, attitudes and practices regarding rabies and the rabies elimination campaign in Bohol as part of the first questionnaire. The results of this questionnaire will be the subject of a later manuscript. The second questionnaire, which is the focus of this article, was designed to collect information on individual dogs within dog-owning households and to assess dog density and dog rabies vaccination coverage. We collected age, sex, vaccination status, vaccination date and primary function on every dog within the sampled household, regardless of how many dogs were present in the household. In addition, we asked about confinement of the dog, how and where the dog was acquired and whether male dogs had been castrated.

Regarding vaccination status, we first asked respondents whether the dog had ever been vaccinated. If the respondent answered, 'yes', we then asked the date (month and year) of the dog's most recent vaccination. Finally, we asked respondents to present proof of vaccination (certificate or collar). Although few respondents were able to produce either their dog's certificate or collar, we felt that most respondents accurately reported their dog's

vaccination status. Thus, we chose to classify dogs as 'vaccinated' if they were reported as having received at least one rabies vaccination in their lifetime, regardless of whether the respondent could present proof of vaccination.

Data analysis

We utilized Intercooled STATA Version 10 (Stata Corporation, College Station, TX, USA) for all statistical analyses. Data were analysed using the survey commands in STATA. Multivariate logistic regression analysis was used to determine odds ratios and assess the potential associations between dog vaccination status and several household demographic variables (employment status of head of household, whether the household had children <15 years of age and whether anyone in the household had ever known someone with rabies) as well as the dog variables of interest from the second questionnaire. For the purposes of the univariate and multivariate analyses, we excluded puppies <3 months of age ($n = 8$) because none of these dogs were vaccinated, and due to the timing of the last vaccination campaign, it is unlikely they would have had the opportunity to be vaccinated.

An odds ratio was considered significant at the 0.05 significance level. For the purposes of constructing a multivariate model, we included any variable of interest that had a significance level of 0.25 in the univariate analyses (Hosmer and Lemeshow, 2000). Variables were then eliminated from the model using a stepwise backward approach. If removing the variable changed the odds ratios of any of the other point estimates in the model by 10% or more, that variable was retained in the final model.

Results

The number of clusters (barangays) selected within each municipality using the modified EPI with PPS sampling design is shown in Fig. 1. Looking at this figure, it appears that the sampling method worked well as municipalities with larger populations typically had a greater number of clusters selected, while those municipalities with smaller populations were less likely to be selected for inclusion in the study.

We interviewed a total of 460 households representing 37 of the 48 municipalities in Bohol and collected information on 541 dogs residing in these households. Overall, dog ownership is common in Bohol. We found that 354 (77%) of the surveyed households reported owning at least one dog, and 118 (33%) of dog-owning households owned more than one dog. The mean number of dogs owned was 1.6 (SD \pm 1.1). For every one dog, there were




Fig. 1. Population (2007) and number of clusters selected using EPI sampling method with population proportional to size (PPS) by municipality, Bohol, Philippines, 2009 [Map was made using the Epi Map feature in Epi Info (Epi Info™ Version 3.5.1; The Centres for Disease Control and Prevention, Atlanta, GA, USA) and with Bohol shape files downloaded from the Natural Resources Database (<http://www.nrdp.co.uk/>)].

approximately four humans. Assuming this ratio holds true throughout the island, we would expect the owned-dog population of Bohol to be approximately 299 297.

Vaccination status

We collected data on vaccination status for 539 of the 541 dogs in our sample. A total of 381 (71%) of the 539 dogs in our study were reported to have been vaccinated for rabies at some time in their lives. However, when respondents were asked when the dog was vaccinated, only 344 (64%) of dogs were reported recently vaccinated in 2008 or 2009. In addition, owners were able to show either the vaccination/registration certificate or collar for only 40% of dogs that were reported vaccinated. Twenty-nine per cent of dogs had never been vaccinated in their lifetime. Table 1 shows the most common reasons given for not vaccinating.

The local government was the most common source of vaccination (65% of dogs). Only 3% of dogs were vaccinated by a private veterinarian. Most respondents had paid some amount of money for vaccination of the dog (88%). The mean amount paid among respondents who reported paying for the vaccination was 75.49 Philippine pesos (approximately 1.74 USD).

Age structure

The dogs in our sample were young, typically around 2 years of age (median = 24.0 months; range = 1–240). Vaccinated dogs were older (median = 36.0 months; range = 3–240) than unvaccinated dogs (median = 12 -

Table 1. Primary reason why dogs have never been vaccinated

	Unvaccinated dogs (n = 158) n (%)
Either respondent or dog not there when vaccine was offered	32 (20)
Dog was believed to be too young	28 (18)
Unable to restrain dog	22 (14)
Dog acquired after vaccination campaign	14 (9)
Vaccine too expensive or no money to pay	12 (8)
Fear of injury or adverse reaction to vaccine	3 (2)
Could not travel to vaccine centre	2 (1)
Unaware of campaign	2 (1)
Other	13 (8)

months; range = 1–216). Puppies <3 months of age were virtually never vaccinated, which is most likely due to the timing of the last vaccination campaign. However, we observed that only 43% of dogs between the ages of 3–11 months of age had ever been vaccinated compared to 73% of those between the ages of 1–2 years. In addition, dogs <1 year of age made up almost 16% of the entire dog population in Bohol. Figure 2 shows vaccination status by age group.

Sex characteristics

Male dogs were preferred to female dogs in Bohol and were about twice as abundant. We found a female-to-male sex ratio of 1 : 2. Males were also more frequently reported to be vaccinated than females, 74% versus 64%, respectively.




Fig. 2. Vaccination status by age group as percentage of all dogs ($n = 539$; Data on age were missing for 118 dogs), Bohol, 2009.




Fig. 3. Age distribution by sex as percentage of all dogs ($n = 539$; Data on age were missing for 118 dogs), Bohol, 2009.

Male dogs were also older (median = 35.0 months; range = 2–240) than female dogs (median = 22.5 months; range = 1–144). Figure 3 shows the age and sex composition of dogs included in our survey.

Reproduction

Approximately 33% of male dogs were castrated in Bohol, representing 22% of all dogs. Vaccinated dogs were significantly more likely to be castrated male dogs ($P < 0.001$). We did not ask about female sterilization, because it is very rare that a female dog has been spayed, mainly due to the cost involved. We did not observe any significant association between vaccination and pregnant female dogs. We also collected information on the number and fate of puppies produced by female dogs in our survey.

Table 2. Reproductive characteristics by vaccination status of the 539 dogs included in the survey

	All dogs ($n = 539$) n (%)	Vaccinated dogs ($n = 381$) n (%)	Unvaccinated dogs ($n = 158$) n (%)	χ^2 (P -value)
Sex				
Male	358 (66)	265 (70)	92 (58)	5.98 (0.01)
Female	181 (34)	116 (30)	65 (41)	
Castrated male				
Yes	117 (22)	106 (25)	11 (6)	22.67 (<0.001)
No	222 (41)	149 (34)	73 (41)	
Pregnant female				
Yes	24 (4)	16 (4)	8 (4)	0.06 (0.80)
No	111 (21)	71 (16)	40 (22)	

The median number of litters produced in the past year was 2 with a range of 1–10. The median number of puppies delivered in the last whelping was 4 with a range of 1–8. However, the median number of puppies surviving beyond 1 month of age was only 2 with a range of 0–7. Table 2 shows reproductive status by vaccination.

Dog acquisition, purpose and management

Most dogs were acquired as gifts (61%) and functioned as guard dogs (75%). A majority of dog owners reported that their dog was confined both night and day (67%). Any confinement (day, night or both day and night) of a dog was significantly associated with vaccination ($P = 0.001$, $P = 0.005$ and $P = 0.01$, respectively). An additional 5% of dogs were reported to be leashed both day and night. However, few dogs (29%) were reported by the interviewer to reside at a residence that had a fence capable of restraining the dog. Table 3 shows the dog characteristics by vaccination status.

Multivariate model

The results of our univariate and multivariate analyses are summarized in Table 4. We found that the dog’s age was positively associated with vaccination, and this finding was statistically significant, even after eliminating dogs <3 months of age from our analyses. Compared to dogs between the ages of 3 and 11 months, the odds of vaccination were approximately 4.6 times greater for dogs between the ages of 12 and 23 months ($P = 0.002$) and approximately 11.7 times greater for dogs between the ages of 36 and 47 months ($P = 0.002$). We also found that dogs had approximately twice the odds of being vaccinated if they were confined both day and night to the household premises, and this result approached statistical significance ($P = 0.07$) in the multivariate model.

Table 3. Dog characteristics by vaccination status of the 539 dogs included in the survey

	All dogs (<i>n</i> = 539) <i>n</i> (%)	Vaccinated dogs (<i>n</i> = 381) <i>n</i> (%)	Unvaccinated dogs (<i>n</i> = 158) <i>n</i> (%)	χ^2 (<i>P</i> -value)
How was dog acquired				(0.13) ^a
Gift	327 (61)	225 (59)	102 (65)	
Purchased	67 (12)	54 (14)	13 (8)	
Born in household	112 (21)	78 (20)	34 (22)	
Found	11 (2)	7 (2)	4 (3)	
Other	13 (2)	12 (3)	1 (1)	
Owner confines dog during day				11.00 (0.001)
Yes	388 (72)	290 (76)	98 (62)	
No	151 (28)	91 (24)	60 (38)	
Owner confines dog during night				7.83 (0.005)
Yes	374 (69)	278 (73)	96 (61)	
No	165 (31)	103 (27)	62 (39)	
Owner confines dog during the day and night				6.34 (0.01)
Yes	360 (67)	267 (70)	93 (59)	
No	179 (33)	114 (30)	65 (41)	
Dog's function				11.00 (0.004)
Guard	394 (75)	273 (73)	121 (80)	
Pet/companion	54 (10)	34 (9)	20 (13)	
Both guard and pet	79 (15)	68 (18)	11 (7)	
Dog resides in multiple dog household				1.09 (0.30)
Yes	312 (58)	226 (59)	86 (54)	
No	227 (42)	155 (41)	72 (46)	

^aFisher's exact test *P*-value.

Female sex appeared to be associated with lower vaccination coverage in the univariate analysis; however, this effect disappeared when age was controlled for in the model. While females are less often vaccinated, this is most likely due to the fact that females are generally younger than males, and younger dogs are less often vaccinated. Odds of vaccination also increased if the owner was employed (OR = 2.1; *P* = 0.07) and decreased if there was a child under the age of 15 in the household (OR = 0.6; *P* = 0.11); however, only owner employment approached statistical significance in the final multivariate model.

Discussion

Dog ecology and demography studies are essential to the planning and long-term success of dog rabies vaccination and disease elimination campaigns (WHO, 1992). With this in mind, we have attempted to gain insight into the owned-dog population of Bohol. The prevalence of dog ownership is high in Bohol, with 77% of sampled households owning at least one dog. In contrast, in the United States, approximately 37% of households own one or more dogs (AVMA, 2007). Previous studies in the Philippines have reported household dog ownership ranging from 40% to 69% (Robinson et al., 1996; Estrada et al.,

2001). However, the mean number of dogs owned in Bohol (approximately 1.6) is similar to that of the United States (1.7; AVMA, 2007) and lower than that found in many other developing countries (Beran, 1982; Fishbein et al., 1992; Robinson et al., 1996; Kitale et al., 2001; Awoyomi et al., 2007; Knobel et al., 2008).

As with most developing countries, a large proportion of the owned-dog population in Bohol is young, with 16% of the dog population <1 year of age and a median age of approximately 2 years. However, compared to some countries in Africa and Asia, the median age of dogs in Bohol was higher. De Balogh et al. (1993) reported a median age of 1 year for dogs residing in a semi-rural village in Zambia (urban dogs in Zambia had a median age of 2 years). Kitale et al. (2001) found that dogs in Kenya had a mean age of only 1.9 years. Even among the studies conducted in the Philippines, dogs in Bohol, on average, lived considerably longer. Estrada et al. (2001) found that female dogs in Mindoro, La Union, Philippines, had a mean age of 1.7 years, while the mean age of male dogs was only 0.4 years. Similarly, Robinson et al. (1996) found a median age of 1 year for dogs in Sorsogon Province, Philippines. Although dogs in Bohol were not as long-lived as dogs in the Coquimbo region of Chile with a median age of 3.0 years (Acosta-Jamett et al., 2010) nor those in Thungsong District, Thailand (Kongkaew et al.,

Table 4. Results of univariate and multivariate logistic regression of factors associated with rabies vaccination in dogs in Bohol^a

Variable	Univariate OR (95% CI)	P-value	Multivariate OR (95% CI)	P-value
HH employed				
Yes	2.1 (1.3–3.6)	0.01	2.1 (0.9–4.8)	0.07
No	Referent	Referent	Referent	Referent
Child under 15				
Yes	0.5 (0.3–0.9)	0.02	0.6 (0.3–1.1)	0.1
No	Referent	Referent	Referent	Referent
Ever known someone with rabies				
Yes	1.1 (0.6–2.0)	0.8		
No	Referent	Referent		
Dog sex				
Female	0.7 (0.4–1.0)	0.01	1.0 (0.5–1.9)	0.9
Male	Referent	Referent	Referent	Referent
Dog age				
3–11 months	Referent	Referent	Referent	Referent
12–23 months	3.7 (1.7–8.3)	0.002	4.6 (1.8–12.0)	0.002
24–35 months	3.4 (1.4–8.1)	0.01	4.5 (1.6–12.9)	0.006
36–47 months	10.7 (3.5–32.7)	<0.001	11.7 (2.5–53.4)	0.002
48+ months	9.4 (4.4–20.3)	<0.001	26.8 (3.9–186.1)	0.001
Dog resides in a multiple dog household				
Yes	1.2 (0.8–1.9)	0.30		
No	Referent	Referent		
Distance from OPV	–0.003 ^b (–0.01 to 0.01)	0.53		
How dog was acquired				
Found	Referent	Referent		
Gift	1.3 (0.4–4.3)	0.71		
Purchased	2.4 (0.5–11.1)	0.26		
Born in household	1.3 (0.4–4.3)	0.65		
Dog came from different municipality				
Yes	1.8 (0.8–3.8)	0.15		
No	Referent	Referent		
Months in current household	0.03 ^b (0.01–0.04)	<0.001	–0.004 ^b (–0.03 to 0.018)	0.72
Dog function				
Guard	Referent	Referent		
Pet	0.8 (0.3–1.7)	0.49		
Both pet and guard	2.7 (1.1–6.6)	0.03		
Owner confines dog day and night				
Yes	1.6 (0.9–3.2)	0.14	2.1 (0.9–4.9)	0.07
No	Referent	Referent	Referent	Referent

^aExcluding eight dogs <3 months of age and two dogs where vaccination status was unknown; *n* = 531.

^bBeta coefficient.

2004), where the mean age was 2.6 years, and compared to dogs in the developed world, lifespan is much shorter among the dogs of Bohol. Owned dogs in the United States have a mean age of approximately 4.5 years (WHO, 1987), while a survey of dogs in Britain reported that 11 years was the approximate mean age at death (Michell, 1999). Aside from the issue of dog welfare, the primary problem with a young dog population is that they are less likely to be vaccinated (Flores-Ibarra and Estrella-Valenzuela, 2004; Suzuki et al., 2008; Kaare et al., 2009), and a younger dog population increases the need for more frequent vaccination campaigns (WHO, 1987). Not surprisingly, we found that the odds of dog vaccination in our

sample increased dramatically with increasing age, and this finding was highly significant.

We determined the sex ratio of male-to-female dogs in Bohol is 2 : 1. This trend has been observed repeatedly in developing countries (Beran and Frith, 1988; Brooks, 1990; Kongkaew et al., 2004; Acosta-Jamett et al., 2010). Male dogs are preferred because the primary function of the dog in Bohol is guarding. Because castration of male dogs is cheaper and more readily available than spaying of female dogs, this is not necessarily a bad trend. However, intact male dogs are often implicated in the spread of rabies due to their inclination towards roaming and fighting (WHO, 2005a); therefore, it is essential that male

dogs be castrated. We found that about a third of male dogs, representing 22% of all dogs in Bohol, are castrated. This is higher than some other developing countries such as Chile (3%; Acosta-Jamett et al., 2010), Zimbabwe (16%; Butler and Bingham, 2000) and Mexico (5%; Flores-Ibarra and Estrella-Valenzuela, 2004; Ortega Pacheco et al., 2007) but lower than Thailand (37%; Kongkaew et al., 2004) and Bolivia (33%; Suzuki et al., 2008).

We found that rabies vaccination coverage in dogs in Bohol was approximately 71%, although only 64% reported having had their dog vaccinated within the previous year (2008 or 2009). It is possible that respondents were simply unaware of when their dog was last vaccinated; however, because few could actually produce a registration card and dogs were seldom seen wearing their registration collars, it was not possible to verify whether or when dogs were actually vaccinated, and this was one of the limitations of our study.

Dogs in our survey were most often not vaccinated due to the unavailability of the dog or the owner at the time vaccination was offered. A small number of dogs were not vaccinated due to an inability to restrain them, and the odds of vaccination were 2.1 times greater in dogs that were confined day and night. In addition, there is still a belief that young dogs cannot be vaccinated, despite the WHO recommendation that age should not be a limiting factor in vaccination (WHO, 1992). Interestingly, we did not find that the fee charged for vaccination/registration prevented many dogs from being vaccinated.

Conclusion

Based on our results, we feel that the programme can be improved in several ways. A more foolproof system for keeping track of vaccinated (registered) dogs needs to be explored further as the identification of vaccinated dogs is crucial to programme evaluation. Offering vaccinations on multiple days at varying times in the same location will help reduce the number of dogs missed by a single vaccination clinic as will encouraging residents to keep their dogs confined or restrained, especially prior to the vaccination campaign. Confinement of a dog may also improve its docility as well as encourage healthy interaction between humans and dogs. Finally, additional public education is needed to counteract the belief that young dogs cannot be vaccinated.

If Bohol and the Philippine governments are able to maintain their level of commitment to controlling rabies, it is clear that the long-term success of the programme is on track. It is also clear that maintaining vaccination coverage of at least 70% in the dog population can prevent rabies in humans.

Due to their intense and successful efforts, the Bohol Rabies Elimination Program won the highly competitive Galing Pook award in 2011 in recognition of the achievements of local government units (Global Alliance for Rabies Control, 2011). As long as the government is able to receive continued support to sustain these efforts, both the dog and human population of Bohol will be protected from death from rabies.

Acknowledgements

The authors wish to thank Asha Kapadia and Sally Vernon with the University of Texas Health Science Center at Houston, School of Public Health (UTSPH), for their contribution to this manuscript as well as UTSPH students Rebecca Bryson, Kara Chappell, Emily Herrington and Melissa Nolan for their tireless support during data collection. In addition, thank you to the staff and volunteers of the Office of the Provincial Veterinarian in Tagbilaran City, Bohol, who made this work possible.

Funding

Dr Lapiz and Dr Miranda received support from the Global Alliance for Rabies Control. Dr Davlin was a PhD student during data collection and development of the original manuscript. She is now a National Research Service Award (NRSA) postdoctoral fellow supported through an institutional training grant from the Eunice Kennedy Shriver National Institute of Child Health & Human Development of the National Institutes of Health (NRSA T32HD055163, Berenson). The content is solely the responsibility of the authors and does not necessarily represent the official views of the Global Alliance for Rabies Control, NICHD, or the NIH.

References

- Acosta-Jamett, G., S. Cleaveland, A. A. Cunningham, and B. M. Bronsvoort, 2010: Demography of domestic dogs in rural and urban areas of the Coquimbo region of Chile and implications for disease transmission. *Prev. Vet. Med.* 94, 272–281.
- American Veterinary Medical Association, 2007: U.S. Pet Ownership & Demographics Sourcebook, 2007 edn. AVMA, Schaumburg, IL.
- Awoyomi, O. J., F. S. Adeyemi, and F. S. Awoyomi, 2007: Socioeconomic factors associated with non-vaccination of dogs against rabies in Ibadan, Nigeria. *Nig. Vet. J.* 28, 59–63.
- Baer, G. M., 2007: The history of rabies. In: Jackson, A. C., and W. H. Wunner (eds), *Rabies*, 2nd edn, pp. 1. Elsevier/Academic Press, Amsterdam; Boston, MA.

- Bennett, S., T. Woods, W. M. Liyanage, and D. L. Smith, 1991: A simplified general method for cluster sample surveys of health in developing countries. *World Health Stat. Q.* 44, 98–106.
- Beran, G. W., 1982: Ecology of dogs in the Central Philippines in relation to rabies control efforts. *Comp. Immunol. Microbiol. Infect. Dis.* 5, 265–270.
- Beran, G. W., and M. Frith, 1988: Domestic animal rabies control: an overview. *Rev. Infect. Dis.* 10(Suppl. 4), S672–S677.
- Brooks, R., 1990: Survey of the dog population of Zimbabwe and its level of rabies vaccination. *Vet. Rec.* 127, 592–596.
- Butler, J. R., and J. Bingham, 2000: Demography and dog-human relationships of the dog population in Zimbabwean communal lands. *Vet. Rec.* 147, 442–446.
- Coleman, P. G., and C. Dye, 1996: Immunization coverage required to prevent outbreaks of dog rabies. *Vaccine* 14, 185–186.
- De Balogh, K. M., A. I. Wandeler, and F. X. Meslin, 1993: A dog ecology study in an urban and a semi-rural area of Zambia. *Onderstepoort J. Vet. Res.* 60, 437–443.
- Dodet, B., 2010: Report of the sixth Asian Rabies Expert Bureau (AREB) meeting, Manila, The Philippines, 10–12 November 2009. *Vaccine* 28, 3265–3268.
- Estrada, R., A. Vos, R. De Leon, and T. Mueller, 2001: Field trial with oral vaccination of dogs against rabies in the Philippines. *BMC Infect. Dis.* 1, 23.
- Fishbein, D. B., N. J. Miranda, P. Merrill, R. A. Camba, M. Meltzer, E. T. Carlos, C. F. Bautista, P. V. Sopungco, L. C. Mangahas, L. M. Hernandez, M. M. Leoncio, D. Mercado, S. Gregorio, E. Salva, J. G. Dobbins, and W. G. Winkler, 1991: Rabies control in the Republic of the Philippines: benefits and costs of elimination. *Vaccine* 9, 581–587.
- Fishbein, D. B., M. G. Frontini, J. G. Dobbins, E. F. Collins, G. Q. Huerta, J. G. Rodriguez, B. Woo-Ming, J. G. Ramos, A. J. Belotto, J. B. Torres, K. M. Yenne, S. B. Linhart, and G. M. Baer, 1992: Prevention of canine rabies in rural Mexico: an epidemiologic study of vaccination campaigns. *Am. J. Trop. Med. Hyg.* 47, 317–327.
- Flores-Ibarra, M., and G. Estrella-Valenzuela, 2004: Canine ecology and socioeconomic factors associated with dogs unvaccinated against rabies in a Mexican city across the US-Mexico border. *Prev. Vet. Med.* 62, 79–87.
- Global Alliance for Rabies Control, 2011: Project Reports, viewed 2012 March 6. Available at: <http://rabiescontrol.net/what-we-do/project-reports.html>.
- Green, S. J., R. D. Alexander, A. M. Gulayan, C. C. Migrino III, J. Jarantill-Paler, and C. A. Courtney, 2002: Bohol Island: Its Coastal Environment Profile. Bohol Environment Management Office, Bohol and Coastal Resource Management Project, Cebu City, Philippines.
- Henderson, R. H., and T. Sundaresan, 1982: Cluster sampling to assess immunization coverage: a review of experience with a simplified sampling method. *Bull. World Health Organ.* 60, 253–260.
- Hosmer, D. W., and S. Lemeshow, 2000: Applied Logistic Regression, 2nd edn. John Wiley & Sons, Inc., New York.
- Kaare, M., T. Lembo, K. Hampson, E. Ernest, A. Estes, C. Mentzel, and S. Cleaveland, 2009: Rabies control in rural Africa: evaluating strategies for effective domestic dog vaccination. *Vaccine* 27, 152–160.
- Kitala, P., J. McDermott, M. Kyule, J. Gathuma, B. Perry, and A. Wandeler, 2001: Dog ecology and demography information to support the planning of rabies control in Machakos District, Kenya. *Acta Trop.* 78, 217–230.
- Knobel, D. L., S. Cleaveland, P. G. Coleman, E. M. Fevre, M. I. Meltzer, M. E. G. Miranda, A. Shaw, J. Zinsstag, and F. X. Meslin, 2005: Re-evaluating the burden of rabies in Africa and Asia. *Bull. World Health Organ.* 83, 360–368.
- Knobel, D. L., M. K. Laurenson, R. R. Kazwala, L. A. Boden, and S. Cleaveland, 2008: A cross-sectional study of factors associated with dog ownership in Tanzania. *BMC Vet. Res.* 4, 5.
- Kongkaew, W., P. Coleman, D. U. Pfeiffer, C. Antarasena, and A. Thiptara, 2004: Vaccination coverage and epidemiological parameters of the owned-dog population in Thungsong District, Thailand. *Prev. Vet. Med.* 65, 105–115.
- Lemeshow, S., and D. Robinson, 1985: Surveys to measure programme coverage and impact: a review of the methodology used by the expanded programme on immunization. *World Health Stat. Q.* 38, 65–75.
- Macpherson, C. N. L., F. Meslin, and A. I. Wandeler (eds), 2000: Dogs, Zoonoses, and Public Health. CABI Pub., New York.
- Michell, A. R., 1999: Longevity of British breeds of dog and its relationships with sex, size, cardiovascular variables and disease. *Vet. Rec.* 145, 625–629.
- Ortega Pacheco, A., J. C. Rodriguez Buenfil, M. E. Bolio Gonzalez, C. H. Sauri Arceo, M. Jimenez Coello, and C. L. Forsberg, 2007: A survey of dog populations in urban and rural areas of Yucatan, Mexico. *Anthrozoos* 20, 261–274.
- Republic of the Philippines-Department of Health, 2011: National rabies prevention and control program. Available at: <http://dev1.doh.gov.ph/content/national-rabies-prevention-and-control-program> (accessed on 14 May 2012).
- Robinson, L. E., M. E. Miranda, N. L. Miranda, and J. E. Childs, 1996: Evaluation of a canine rabies vaccination campaign and characterization of owned-dog populations in the Philippines. *Southeast Asian J. Trop. Med. Public Health* 27, 250–256.
- Rupprecht, C. E., J. Barrett, D. Briggs, F. Cliquet, A. R. Fooks, B. Lumlerdacha, F. X. Meslin, T. Muler, L. H. Nel, C. Schneider, N. Tordo, and A. I. Wandeler, 2008: Can rabies be eradicated? *Dev. Biol. (Basel)* 131, 95–122.
- Suzuki, K., J. A. Pereira, L. A. Frias, R. Lopez, L. E. Mutinelli, and E. R. Pons, 2008: Rabies-vaccination coverage and profiles of the owned-dog population in Santa Cruz de la Sierra, Bolivia. *Zoonoses Public Health* 55, 177–183.

- WHO, 1987: Guidelines for Dog Rabies Control. World Health Organization, Geneva, Switzerland.
- WHO, 1992: Expert Committee on Rabies: Eighth Report. World Health Organization, Geneva, Switzerland.
- WHO, 2002: Strategies for the Control and Elimination of Rabies in Asia: Report of a WHO Interregional Consultation. World Health Organization Department of Epidemic and Pandemic Alert and Response, Geneva, Switzerland.
- WHO, 2005a: WHO Expert Consultation on Rabies. First Report. World Health Organization, Geneva, Switzerland.
- WHO, 2005b: Immunization Coverage Cluster Survey – Reference Manual. Department of Immunization, Vaccines and Biologicals, Family and Community Health, World Health Organization, Geneva, Switzerland.